

THE EQUIVALENCE BETWEEN A NEW MULTISTEP ITERATION, S-ITERATION AND SOME OTHER ITERATIVE SCHEMES

FAIK GÜRSOY, VATAN KARAKAYA, AND B. E. RHOADES

ABSTRACT. In this paper, we show that Picard, Krasnoselskij, Mann, Ishikawa, new two step, Noor, multistep, new multistep, SP and S-iterative schemes are equivalent for contractive-like operators.

1. INTRODUCTION AND PRELIMINARIES

In the last four decades, attention of researchers has been focused on the introduction and the convergences of various iteration procedures, e.g. see [7, 14, 18, 21, 30, 26, 16, 11], among others, for approximate fixed points of certain classes of self-operators. During the past 11 years, a large literature has developed around theme establishing equivalence between convergences of some well-known iterative schemes deal with various classes of operators. The authors who have made contributions to the study of equivalence between various iterative schemes are Rhoades and Şoltuz [[1]-[8]], Berinde [28], Şoltuz [22, 23], Olaleru and Akewe [13], Chang et al [20] and several references therein.

The aim of this paper is to show equivalence between convergences of a new multistep iteration, which is unifies and developes the iterative algorithms presented in [26] and [16], S-iteration and some other iterative schemes.

As a background of our exposition, we now mention some contractive mappings and iteration schemes.

In [27] Zamfirescu established an important generalization of the Banach fixed point theorem using the following contractive condition: For a mapping $T : E \rightarrow E$, there exist real numbers a, b, c satisfying $0 < a < 1$, $0 < b, c < 1/2$ such that, for each pair $x, y \in X$, at least one of the following is true:

$$(1.1) \quad \begin{cases} (z_1) & \|Tx - Ty\| \leq a \|x - y\|, \\ (z_2) & \|Tx - Ty\| \leq b (\|x - Tx\| + \|y - Ty\|), \\ (z_3) & \|Tx - Ty\| \leq c (\|x - Ty\| + \|y - Tx\|). \end{cases}$$

A mapping T satisfying the contractive conditions (z_1) , (z_2) and (z_3) in (1.1) is called a Zamfirescu operator. An operator satisfying condition (z_2) is called a *Kannan operator*, while the mapping satisfying condition (z_3) is called a *Chatterjea operator*. As shown in [29], the contractive condition (1.1) leads to

$$(1.2) \quad \begin{cases} (b_1) & \|Tx - Ty\| \leq \delta \|x - y\| + 2\delta \|x - Tx\| \text{ if one use } (z_2), \\ \text{and} & \\ (b_2) & \|Tx - Ty\| \leq \delta \|x - y\| + 2\delta \|x - Ty\| \text{ if one use } (z_3), \end{cases}$$

2000 *Mathematics Subject Classification.* Primary 47H10.

Key words and phrases. New multistep iteration, S-iteration, Equivalence of iterations, Contractive-like operator.

for all $x, y \in E$ where $\delta := \max \left\{ a, \frac{b}{1-b}, \frac{c}{1-c} \right\}$, $\delta \in [0, 1)$, and it was shown that this class of operators is wider than the class of Zamfirescu operators. Any mapping satisfying condition (b₁) or (b₂) is called a quasi-contractive operator.

Extending the above definition, Osilike and Udomene [15] considered operators T for which there exist real numbers $L \geq 0$ and $\delta \in [0, 1)$ such that for all $x, y \in E$,

$$(1.3) \quad \|Tx - Ty\| \leq \delta \|x - y\| + L \|x - Tx\|.$$

Imoru and Olantiwo [9] gave a more general definition: The operator T is called a contractive-like operator if there exists a constant $\delta \in [0, 1)$ and a strictly increasing and continuous function $\varphi : [0, \infty) \rightarrow [0, \infty)$ with $\varphi(0) = 0$, such that, for each $x, y \in E$,

$$(1.4) \quad \|Tx - Ty\| \leq \delta \|x - y\| + \varphi(\|x - Tx\|).$$

A map satisfying (1.4) need not have a fixed point, even if E is complete. For example, let $E = [0, \infty)$, and define T by

$$Tx = \begin{cases} 1.0, & 0 \leq x \leq 0.8, \\ 0.6, & 0.8 < x. \end{cases}$$

Without loss of generality we may assume that $x < y$. Then, for $0 \leq x < y \leq 0.8$ or $0.8 < x < y$, $\|Tx - Ty\| = 0$, and (1.4) automatically satisfied.

If $0 \leq x \leq 0.8 < y$, then $\|Tx - Ty\| = 0.4$.

Define φ by $\varphi(t) = Lt$ for any $L \geq 2$. Then φ is increasing, continuous, and $\varphi(0) = 0$. Also, $\|x - Tx\| = 1 - x$, so that $\varphi(\|x - Tx\|) = L(1 - x) \geq 0.2L \geq 0.4$.

Therefore

$$0.4 = \|Tx - Ty\| \leq L \|x - Tx\| \leq \delta \|x - y\| + L \|x - Tx\|$$

for any $0 \leq \delta < 1$, and (1.4) is satisfied for $0 \leq x \leq 0.8 < y$. But T has no fixed point.

However, using (1.4) it is obvious that, if T has a fixed point, then it is unique.

Throughout this paper \mathbb{N} denotes the set of all nonnegative integers. Let X be a Banach space and $E \subset X$ a nonempty closed, convex subset of X , and T a self map on E . Define $F_T := \{p \in X : p = Tp\}$ to be the set of fixed points of T . Let $\{\alpha_n\}_{n=0}^\infty$, $\{\beta_n\}_{n=0}^\infty$, $\{\gamma_n\}_{n=0}^\infty$ and $\{\beta_n^i\}_{n=0}^\infty$, $i = \overline{1, k-2}$, $k \geq 2$ be real sequences in $[0, 1)$ satisfying certain conditions.

It is well known that Picard iteration procedure is defined by

$$(1.5) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = Tx_n, \quad n \in \mathbb{N}. \end{cases}$$

The Mann iterative scheme [30] is defined by

$$(1.6) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n Tx_n, \quad n \in \mathbb{N}. \end{cases}$$

Taking $\alpha_n = \lambda$ (constant) in (1.6), we get Krasnoselskij iteration procedure as follows

$$(1.7) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \lambda)x_n + \lambda Tx_n, \quad n \in \mathbb{N}. \end{cases}$$

A sequence $\{x_n\}_{n=0}^{\infty}$ defined by

$$(1.8) \quad \begin{cases} x_0 \in E, \\ y_n = (1 - \beta_n)x_n + \beta_n T x_n, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n, \end{cases} \quad n \in \mathbb{N},$$

is commonly known as the Ishikawa iterative method [21].

The following iteration scheme introduced by Noor [14]

$$(1.9) \quad \begin{cases} x_0 \in E, \\ z_n = (1 - \gamma_n)x_n + \gamma_n T x_n, \\ y_n = (1 - \beta_n)x_n + \beta_n T z_n, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n, \end{cases} \quad n \in \mathbb{N}.$$

In year 2004, Rhoades and Soltuz introduced in [7] a multistep procedure defined by

$$(1.10) \quad \begin{cases} x_0 \in E, \\ y_n^{k-1} = (1 - \beta_n^{k-1})x_n + \beta_n^{k-1}T x_n, \quad k \geq 2, \\ y_n^i = (1 - \beta_n^i)x_n + \beta_n^i T y_n^{i+1}, \quad i = \overline{1, k-2}, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n^1, \end{cases} \quad n \in \mathbb{N}.$$

The iteration processes (1.5), (1.6), (1.7), (1.8) and (1.9) can be viewed as the special cases of the iteration procedure (1.10).

The sequence $\{x_n\}_{n=0}^{\infty}$ defined by

$$(1.11) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)T x_n + \alpha_n T y_n, \\ y_n = (1 - \beta_n)x_n + \beta_n T x_n, \end{cases} \quad n \in \mathbb{N}$$

is known as the S-iteration process [10, 17, 18].

In 2008, S.Thianwan [26] defined the following iterative process

$$(1.12) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)y_n + \alpha_n T y_n, \\ y_n = (1 - \beta_n)x_n + \beta_n T x_n, \end{cases} \quad n \in \mathbb{N}.$$

Recently Phuengrattana and Suantai introduced an SP iteration method [16] by

$$(1.13) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)y_n + \alpha_n T y_n, \\ y_n = (1 - \beta_n)z_n + \beta_n T z_n, \\ z_n = (1 - \gamma_n)x_n + \gamma_n T x_n, \end{cases} \quad n \in \mathbb{N}.$$

The following iteration scheme is first employed in [11] as a special case of Jungck multistep-SP iteration [12]: For arbitrary fixed order $k \geq 2$,

$$(1.14) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n)y_n^1 + \alpha_n T y_n^1, \\ y_n^1 = (1 - \beta_n^1)y_n^2 + \beta_n^1 T y_n^2, \\ y_n^2 = (1 - \beta_n^2)y_n^3 + \beta_n^2 T y_n^3, \\ \dots \\ y_n^{k-2} = (1 - \beta_n^{k-2})y_n^{k-1} + \beta_n^{k-2} T y_n^{k-1}, \\ y_n^{k-1} = (1 - \beta_n^{k-1})x_n + \beta_n^{k-1} T x_n, \end{cases} \quad n \in \mathbb{N}.$$

or, in short,

$$(1.15) \quad \begin{cases} x_0 \in E, \\ x_{n+1} = (1 - \alpha_n) y_n^1 + \alpha_n T y_n^1, \\ y_n^i = (1 - \beta_n^i) y_n^{i+1} + \beta_n^i T y_n^{i+1}, \quad i = \overline{1, k-2}, \\ y_n^{k-1} = (1 - \beta_n^{k-1}) x_n + \beta_n^{k-1} T x_n, \quad n \in \mathbb{N}. \end{cases}$$

where, $\{\alpha_n\}_{n=0}^\infty \subset [0, 1)$ is a sequence of real numbers such that,

$$(1.16) \quad \sum_{n=0}^\infty \alpha_n = \infty$$

and

$$(1.17) \quad \{\beta_n^i\}_{n=0}^\infty \subset [0, 1), \quad i = \overline{1, k-1}.$$

Remark 1. If $\gamma_n = 0$, then SP iteration (1.13) reduces to iterative scheme (1.12). By taking $k = 3$ and $k = 2$ in (1.15) we obtain the iterations (1.13) and (1.12), respectively.

The following Lemma will be useful to prove the main results of this work and is important by itself.

Lemma 1. [31] Let $\{a_n\}_{n=0}^\infty$ be a nonnegative sequence which satisfies the following inequality

$$(1.18) \quad a_{n+1} \leq (1 - \mu_n) a_n + \rho_n,$$

where $\mu_n \in (0, 1)$, for all $n \geq n_0$, $\sum_{n=0}^\infty \mu_n = \infty$, and $\rho_n = o(\mu_n)$. Then $\lim_{n \rightarrow \infty} a_n = 0$.

2. MAIN RESULTS

Theorem 1. Let $T : E \rightarrow E$ be an operator satisfying condition (1.4) with $F_T \neq \emptyset$. If $x_0 = u_0 \in E$ and $\alpha_n \geq A > 0, \forall n \in \mathbb{N}$, then the following are equivalent:

- (1) The Mann iteration (1.6) converges to $p \in F_T$,
- (2) The new multistep iteration (1.15) converges to $p \in F_T$.

Proof. Firstly, we start to prove the implication (1) \Rightarrow (2): Suppose that the Mann iteration (1.6) converges to p . Using (1.6), (1.15), and (1.4) we have the following estimates:

$$(2.1) \quad \begin{aligned} \|u_{n+1} - x_{n+1}\| &= \|(1 - \alpha_n)(u_n - y_n^1) + \alpha_n(Tu_n - Ty_n^1)\| \\ &\leq (1 - \alpha_n)\|u_n - y_n^1\| + \alpha_n\|Tu_n - Ty_n^1\| \\ &\leq (1 - \alpha_n)\|u_n - y_n^1\| + \alpha_n\{\delta\|u_n - y_n^1\| + \varphi(\|u_n - Tu_n\|)\} \\ &= [1 - \alpha_n(1 - \delta)]\|u_n - y_n^1\| + \alpha_n\varphi(\|u_n - Tu_n\|), \end{aligned}$$

$$\begin{aligned}
\|u_n - y_n^1\| &= \|u_n - (1 - \beta_n^1) y_n^2 - \beta_n^1 T y_n^2\| \\
&= \|u_n - \beta_n^1 u_n + \beta_n^1 u_n - (1 - \beta_n^1) y_n^2 - \beta_n^1 T y_n^2\| \\
&= \|(1 - \beta_n^1) (u_n - y_n^2) + \beta_n^1 (u_n - T y_n^2)\| \\
&\leq (1 - \beta_n^1) \|u_n - y_n^2\| + \beta_n^1 \|u_n - T y_n^2\| \\
&= (1 - \beta_n^1) \|u_n - y_n^2\| + \beta_n^1 \|u_n - T u_n + T u_n - T y_n^2\| \\
&\leq (1 - \beta_n^1) \|u_n - y_n^2\| + \beta_n^1 \|T u_n - T y_n^2\| + \beta_n^1 \|u_n - T u_n\| \\
&\leq (1 - \beta_n^1) \|u_n - y_n^2\| + \beta_n^1 \delta \|u_n - y_n^2\| + \beta_n^1 \varphi (\|u_n - T u_n\|) \\
&\quad + \beta_n^1 \|u_n - T u_n\| \\
(2.2) \quad &= [1 - \beta_n^1 (1 - \delta)] \|u_n - y_n^2\| + \beta_n^1 \{ \|u_n - T u_n\| + \varphi (\|u_n - T u_n\|) \},
\end{aligned}$$

$$\begin{aligned}
\|u_n - y_n^2\| &= \|(1 - \beta_n^2) (u_n - y_n^3) + \beta_n^2 (u_n - T y_n^3)\| \\
&\leq (1 - \beta_n^2) \|u_n - y_n^3\| + \beta_n^2 \|u_n - T y_n^3\| \\
&\leq (1 - \beta_n^2) \|u_n - y_n^3\| + \beta_n^2 \|T u_n - T y_n^3\| + \beta_n^2 \|u_n - T u_n\| \\
&\leq (1 - \beta_n^2) \|u_n - y_n^3\| + \beta_n^2 \delta \|u_n - y_n^3\| + \beta_n^2 \varphi (\|u_n - T u_n\|) \\
&\quad + \beta_n^2 \|u_n - T u_n\| \\
(2.3) \quad &= [1 - \beta_n^2 (1 - \delta)] \|u_n - y_n^3\| + \beta_n^2 \{ \|u_n - T u_n\| + \varphi (\|u_n - T u_n\|) \},
\end{aligned}$$

$$\begin{aligned}
\|u_n - y_n^3\| &= \|(1 - \beta_n^3) (u_n - y_n^4) + \beta_n^3 (u_n - T y_n^4)\| \\
&\leq (1 - \beta_n^3) \|u_n - y_n^4\| + \beta_n^3 \|u_n - T y_n^4\| \\
&\leq (1 - \beta_n^3) \|u_n - y_n^4\| + \beta_n^3 \|T u_n - T y_n^4\| + \beta_n^3 \|u_n - T u_n\| \\
&\leq (1 - \beta_n^3) \|u_n - y_n^4\| + \beta_n^3 \delta \|u_n - y_n^4\| + \beta_n^3 \varphi (\|u_n - T u_n\|) \\
&\quad + \beta_n^3 \|u_n - T u_n\| \\
(2.4) \quad &= [1 - \beta_n^3 (1 - \delta)] \|u_n - y_n^4\| + \beta_n^3 \{ \|u_n - T u_n\| + \varphi (\|u_n - T u_n\|) \}.
\end{aligned}$$

By combining (2.1), (2.2), (2.3), and (2.4) we obtain

$$\begin{aligned}
\|u_{n+1} - x_{n+1}\| &\leq [1 - \alpha_n (1 - \delta)] [1 - \beta_n^1 (1 - \delta)] [1 - \beta_n^2 (1 - \delta)] \\
&\quad [1 - \beta_n^3 (1 - \delta)] \|u_n - y_n^4\| \\
&\quad + [1 - \alpha_n (1 - \delta)] \{ [1 - \beta_n^1 (1 - \delta)] [1 - \beta_n^2 (1 - \delta)] \beta_n^3 \\
&\quad + [1 - \beta_n^1 (1 - \delta)] \beta_n^2 + \beta_n^1 \} \{ \|u_n - T u_n\| + \varphi (\|u_n - T u_n\|) \} \\
(2.5) \quad &\quad + \alpha_n \varphi (\|u_n - T u_n\|)
\end{aligned}$$

Continuing the above process we have

$$\begin{aligned}
\|u_{n+1} - x_{n+1}\| &\leq [1 - \alpha_n (1 - \delta)] [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-2} (1 - \delta)] \|u_n - y_n^{k-1}\| \\
&\quad + [1 - \alpha_n (1 - \delta)] \left\{ [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-3} (1 - \delta)] \beta_n^{k-2} \right. \\
&\quad + \cdots + [1 - \beta_n^1 (1 - \delta)] \beta_n^2 + \beta_n^1 \} \{ \|u_n - T u_n\| + \varphi (\|u_n - T u_n\|) \} \\
(2.6) \quad &\quad + \alpha_n \varphi (\|u_n - T u_n\|).
\end{aligned}$$

Again using (1.15), and (1.4) we get

$$\begin{aligned}
\|u_n - y_n^{k-1}\| &= \left\| \left(1 - \beta_n^{k-1}\right) (u_n - x_n) + \beta_n^{k-1} (u_n - Tx_n) \right\| \\
&\leq \left(1 - \beta_n^{k-1}\right) \|u_n - x_n\| + \beta_n^{k-1} \|u_n - Tx_n\| \\
&\leq \left(1 - \beta_n^{k-1}\right) \|u_n - x_n\| + \beta_n^{k-1} \|Tu_n - Tx_n\| + \beta_n^{k-1} \|u_n - Tu_n\| \\
(2.7) \quad &\leq \left[1 - \beta_n^{k-1} (1 - \delta)\right] \|u_n - x_n\| + \beta_n^{k-1} \{\|u_n - Tu_n\| + \varphi(\|u_n - Tu_n\|)\}.
\end{aligned}$$

Since $\delta \in [0, 1)$ and $\{\alpha_n\}_{n=0}^\infty, \{\beta_n^i\}_{n=0}^\infty \subset [0, 1)$ for $i = \overline{1, k-1}$, we have

$$(2.8) \quad [1 - \alpha_n (1 - \delta)] [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-1} (1 - \delta)] \leq [1 - \alpha_n (1 - \delta)].$$

Using inequality (2.8) and the assumption $\alpha_n \geq A > 0, \forall n \in \mathbb{N}$ in resultant inequality obtained by substituting (2.7) in (2.6) we get

$$\begin{aligned}
\|u_{n+1} - x_{n+1}\| &\leq [1 - A (1 - \delta)] \|u_n - x_n\| \\
&\quad + [1 - A (1 - \delta)] \left\{ [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-2} (1 - \delta)] \beta_n^{k-1} \right. \\
&\quad \left. + \cdots + [1 - \beta_n^1 (1 - \delta)] \beta_n^2 + \beta_n^1 \right\} \{\|u_n - Tu_n\| + \varphi(\|u_n - Tu_n\|)\} \\
(2.9) \quad &\quad + \alpha_n \varphi(\|u_n - Tu_n\|).
\end{aligned}$$

Define

$$\begin{aligned}
a_n &: = \|u_n - x_n\|, \\
\mu_n &: = A (1 - \delta) \in (0, 1), \\
\rho_n &: = [1 - A (1 - \delta)] \left\{ [1 - \beta_n^1 (1 - \delta)] \cdots [1 - \beta_n^{k-2} (1 - \delta)] \beta_n^{k-1} \right. \\
&\quad \left. + \cdots + [1 - \beta_n^1 (1 - \delta)] \beta_n^2 + \beta_n^1 \right\} \{\|u_n - Tu_n\| + \varphi(\|u_n - Tu_n\|)\} \\
&\quad + \alpha_n \varphi(\|u_n - Tu_n\|).
\end{aligned}$$

Since $\lim_{n \rightarrow \infty} \|u_n - p\| = 0$ and $Tp = p \in F_T$, it follows from (1.4) that

$$\begin{aligned}
0 &\leq \|u_n - Tu_n\| \\
&\leq \|u_n - p\| + \|Tp - Tu_n\| \\
&\leq \|u_n - p\| + \delta \|p - u_n\| + \varphi(\|p - Tp\|) \\
(2.10) \quad &= (1 + \delta) \|u_n - p\| \rightarrow 0 \text{ as } n \rightarrow \infty,
\end{aligned}$$

which implies $\lim_{n \rightarrow \infty} \|u_n - Tu_n\| = 0$; namely $\rho_n = o(\mu_n)$. Hence an application of Lemma 1 to (2.10) yields $\lim_{n \rightarrow \infty} \|u_n - x_n\| = 0$. Since $u_n \rightarrow p$ as $n \rightarrow \infty$ by assumption, we derive

$$(2.11) \quad \|x_n - p\| \leq \|x_n - u_n\| + \|u_n - p\|$$

and this implies that $\lim_{n \rightarrow \infty} x_n = p$.

(2) \Rightarrow (1) : Assume $x_n \rightarrow p$ as $n \rightarrow \infty$. Using (1.6), (1.15) and (1.4), we have the following estimates:

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &= \|(1 - \alpha_n) (y_n^1 - u_n) + \alpha_n (Ty_n^1 - Tu_n)\| \\
&\leq (1 - \alpha_n) \|y_n^1 - u_n\| + \alpha_n \|Ty_n^1 - Tu_n\| \\
&\leq (1 - \alpha_n) \|y_n^1 - u_n\| + \alpha_n \{\delta \|y_n^1 - u_n\| + \varphi(\|y_n^1 - Ty_n^1\|)\} \\
(2.12) \quad &= [1 - \alpha_n (1 - \delta)] \|y_n^1 - u_n\| + \alpha_n \varphi(\|y_n^1 - Ty_n^1\|),
\end{aligned}$$

$$\begin{aligned}
\|y_n^1 - u_n\| &= \|(1 - \beta_n^1) y_n^2 + \beta_n^1 T y_n^2 - u_n\| \\
&= \|(1 - \beta_n^1) y_n^2 + \beta_n^1 T y_n^2 - u_n (1 - \beta_n^1 + \beta_n^1)\| \\
&= \|(1 - \beta_n^1) (y_n^2 - u_n) + \beta_n^1 (T y_n^2 - u_n)\| \\
&\leq (1 - \beta_n^1) \|y_n^2 - u_n\| + \beta_n^1 \|T y_n^2 - u_n\| \\
&\leq (1 - \beta_n^1) \|y_n^2 - u_n\| + \beta_n^1 \|T y_n^2 - y_n^2 + y_n^2 - u_n\| \\
&\leq (1 - \beta_n^1) \|y_n^2 - u_n\| + \beta_n^1 \|y_n^2 - u_n\| + \beta_n^1 \|T y_n^2 - y_n^2\| \\
(2.13) \quad &= \|y_n^2 - u_n\| + \beta_n^1 \|T y_n^2 - y_n^2\|,
\end{aligned}$$

$$\begin{aligned}
\|y_n^2 - u_n\| &= \|(1 - \beta_n^2) y_n^3 + \beta_n^2 T y_n^3 - u_n\| \\
&= \|(1 - \beta_n^2) (y_n^3 - u_n) + \beta_n^2 (T y_n^3 - u_n)\| \\
&\leq (1 - \beta_n^2) \|y_n^3 - u_n\| + \beta_n^2 \|T y_n^3 - u_n\| \\
&\leq (1 - \beta_n^2) \|y_n^3 - u_n\| + \beta_n^2 \|y_n^3 - u_n\| + \beta_n^2 \|T y_n^3 - y_n^3\| \\
(2.14) \quad &= \|y_n^3 - u_n\| + \beta_n^2 \|T y_n^3 - y_n^3\|.
\end{aligned}$$

By combining (2.12), (2.13), and (2.14) we obtain

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &\leq [1 - \alpha_n (1 - \delta)] \|y_n^3 - u_n\| + [1 - \alpha_n (1 - \delta)] \beta_n^2 \|T y_n^3 - y_n^3\| \\
(2.15) \quad &+ [1 - \alpha_n (1 - \delta)] \beta_n^1 \|T y_n^2 - y_n^2\| + \alpha_n \varphi (\|y_n^1 - T y_n^1\|)
\end{aligned}$$

Continuing in a similar way, we have

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &\leq [1 - \alpha_n (1 - \delta)] \|y_n^{k-1} - u_n\| \\
&+ [1 - \alpha_n (1 - \delta)] \beta_n^{k-2} \|T y_n^{k-1} - y_n^{k-1}\| \\
(2.16) \quad &+ \cdots + [1 - \alpha_n (1 - \delta)] \beta_n^1 \|T y_n^2 - y_n^2\| + \alpha_n \varphi (\|y_n^1 - T y_n^1\|)
\end{aligned}$$

Using now (1.15) we have

$$\begin{aligned}
\|y_n^{k-1} - u_n\| &= \|(1 - \beta_n^{k-1}) x_n + \beta_n^{k-1} T x_n - u_n\| \\
&\leq (1 - \beta_n^{k-1}) \|x_n - u_n\| + \beta_n^{k-1} \|T x_n - u_n\| \\
&\leq (1 - \beta_n^{k-1}) \|x_n - u_n\| + \beta_n^{k-1} \|x_n - u_n\| + \beta_n^{k-1} \|T x_n - x_n\| \\
(2.17) \quad &\leq \|x_n - u_n\| + \beta_n^{k-1} \|T x_n - x_n\|.
\end{aligned}$$

Substituting (2.17) in (2.16) and utilizing the assumption $\alpha_n \geq A > 0, \forall n \in \mathbb{N}$ we get

$$\begin{aligned}
\|x_{n+1} - u_{n+1}\| &\leq [1 - A (1 - \delta)] \|x_n - u_n\| \\
&+ [1 - A (1 - \delta)] \left\{ \beta_n^{k-1} \|T x_n - x_n\| + \beta_n^{k-2} \|T y_n^{k-1} - y_n^{k-1}\| \right. \\
(2.18) \quad &+ \cdots + \beta_n^1 \|T y_n^2 - y_n^2\| \left. \right\} + \alpha_n \varphi (\|y_n^1 - T y_n^1\|).
\end{aligned}$$

Now define

$$\begin{aligned}
a_n &: = \|u_n - x_n\|, \\
\mu_n &: = A (1 - \delta) \in (0, 1), \\
\rho_n &: = + [1 - A (1 - \delta)] \left\{ \beta_n^{k-1} \|T x_n - x_n\| + \beta_n^{k-2} \|T y_n^{k-1} - y_n^{k-1}\| \right. \\
&\quad \left. + \cdots + \beta_n^1 \|T y_n^2 - y_n^2\| \right\} + \alpha_n \varphi (\|y_n^1 - T y_n^1\|).
\end{aligned}$$

Since $\lim_{n \rightarrow \infty} \|x_n - p\| = 0$ and $Tp = p \in F_T$, it follows from (1.4) that

$$\begin{aligned}
 0 &\leq \|x_n - Tx_n\| \\
 &\leq \|x_n - p\| + \|Tp - Tx_n\| \\
 &\leq \|x_n - p\| + \delta \|p - x_n\| + \varphi(\|p - Tp\|) \\
 (2.19) \quad &= (1 + \delta) \|x_n - p\| \rightarrow 0 \text{ as } n \rightarrow \infty.
 \end{aligned}$$

Utilizing (1.4), (1.15), and (1.17), we have

$$\begin{aligned}
 0 &\leq \|y_n^1 - Ty_n^1\| = \|y_n^1 - p + p - Ty_n^1\| \\
 &\leq \|y_n^1 - p\| + \|Tp - Ty_n^1\| \\
 &\leq \|y_n^1 - p\| + \delta \|p - y_n^1\| + \varphi(\|p - Tp\|) \\
 &= (1 + \delta) \|y_n^1 - p\| \\
 &= (1 + \delta) \|(1 - \beta_n^1) y_n^2 + \beta_n^1 Ty_n^2 - p(1 - \beta_n^1 + \beta_n^1)\| \\
 &\leq (1 + \delta) \{(1 - \beta_n^1) \|y_n^2 - p\| + \beta_n^1 \|Ty_n^2 - Tp\|\} \\
 &\leq (1 + \delta) \{(1 - \beta_n^1) \|y_n^2 - q\| + \beta_n^1 \delta \|y_n^2 - p\|\} \\
 &= (1 + \delta) [1 - \beta_n^1 (1 - \delta)] \|y_n^2 - p\| \\
 &= (1 + \delta) [1 - \beta_n^1 (1 - \delta)] \|(1 - \beta_n^2) y_n^3 + \beta_n^2 Ty_n^3 - p(1 - \beta_n^2 + \beta_n^2)\| \\
 &\leq (1 + \delta) [1 - \beta_n^1 (1 - \delta)] \{(1 - \beta_n^2) \|y_n^3 - q\| + \beta_n^2 \|Ty_n^3 - Tp\|\} \\
 &\leq (1 + \delta) [1 - \beta_n^1 (1 - \delta)] [1 - \beta_n^2 (1 - \delta)] \|y_n^3 - p\| \\
 &\quad \dots \\
 &\leq (1 + \delta) [1 - \beta_n^1 (1 - \delta)] \dots [1 - \beta_n^{k-2} (1 - \delta)] \|y_n^{k-1} - p\| \\
 &\leq (1 + \delta) [1 - \beta_n^1 (1 - \delta)] \dots [1 - \beta_n^{k-1} (1 - \delta)] \|x_n - p\| \\
 (2.20) \quad &\leq (1 + \delta) \|x_n - p\| \rightarrow 0 \text{ as } n \rightarrow \infty.
 \end{aligned}$$

It is easy to see from (2.20) that this result is also valid for $\|Ty_n^2 - y_n^2\|, \dots, \|Ty_n^{k-1} - y_n^{k-1}\|$.

Since φ is continuous, we have

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \|x_n - Tx_n\| &= \lim_{n \rightarrow \infty} \varphi(\|y_n^1 - Ty_n^1\|) \\
 (2.21) \quad &= \lim_{n \rightarrow \infty} \|y_n^2 - Ty_n^2\| = \dots = \lim_{n \rightarrow \infty} \|y_n^{k-1} - Ty_n^{k-1}\| = 0,
 \end{aligned}$$

that is $\rho_n = o(\mu_n)$. Hence an application of Lemma 1 to (2.18) lead to $\lim_{n \rightarrow \infty} \|x_n - u_n\| = 0$. Since $x_n \rightarrow p$ as $n \rightarrow \infty$ by assumption, we derive

$$(2.22) \quad \|u_n - p\| \leq \|u_n - x_n\| + \|x_n - p\|$$

and this implies that $\lim_{n \rightarrow \infty} u_n = p$. \square

Theorem 2. Let $T : E \rightarrow E$ be an operator satisfying condition (1.4) with $F_T \neq \emptyset$. If $x_0 = u_0 \in E$ and $\alpha_n \geq A > 0, \forall n \in \mathbb{N}$, then the following are equivalent:

- (1) The Mann iteration (1.6) converges to $p \in F_T$,
- (2) The S-iteration (1.11) converges to $p \in F_T$.

Proof. Firstly, we start to prove the implication (1) \Rightarrow (2): Suppose that the Mann iteration (1.6) converges to p . Using (1.4), (1.6), and (1.11) we have the following

estimates:

$$\begin{aligned}
 \|u_{n+1} - x_{n+1}\| &= \|(1 - \alpha_n)(u_n - Tx_n) + \alpha_n(Tu_n - Ty_n)\| \\
 &\leq (1 - \alpha_n)\|u_n - Tx_n\| + \alpha_n\|Tu_n - Ty_n\| \\
 (2.23) \quad &\leq (1 - \alpha_n)\|u_n - Tx_n\| + \alpha_n\delta\|u_n - y_n\| + \alpha_n\varphi(\|u_n - Tu_n\|),
 \end{aligned}$$

$$\begin{aligned}
 \|u_n - y_n\| &= \|u_n - (1 - \beta_n)x_n - \beta_nTx_n\| \\
 &= \|u_n - \beta_nu_n + \beta_nu_n - (1 - \beta_n)x_n - \beta_nTx_n\| \\
 (2.24) \quad &\leq (1 - \beta_n)\|u_n - x_n\| + \beta_n\|u_n - Tx_n\|,
 \end{aligned}$$

$$\begin{aligned}
 \|u_n - Tx_n\| &= \|u_n - Tu_n + Tu_n - Tx_n\| \\
 &\leq \|u_n - Tu_n\| + \|Tu_n - Tx_n\| \\
 (2.25) \quad &\leq \|u_n - Tu_n\| + \delta\|u_n - x_n\| + \varphi(\|u_n - Tu_n\|).
 \end{aligned}$$

By combining (2.23), (2.24), and (2.25) we obtain

$$\begin{aligned}
 \|u_{n+1} - x_{n+1}\| &\leq \{(1 - \alpha_n)\delta + \alpha_n\delta[1 - \beta_n(1 - \delta)]\}\|u_n - x_n\| \\
 &\quad + [1 - \alpha_n + \alpha_n\beta_n\delta]\|u_n - Tu_n\| \\
 (2.26) \quad &\quad + [1 + \alpha_n\beta_n\delta]\varphi(\|u_n - Tu_n\|).
 \end{aligned}$$

Since $\delta, \alpha_n, \beta_n \in [0, 1]$ for all $n \in \mathbb{N}$,

$$(2.27) \quad (1 - \alpha_n)\delta < 1 - \alpha_n, \quad 1 - \beta_n(1 - \delta) < 1.$$

Using (2.27) and the assumption $\alpha_n \geq A > 0, \forall n \in \mathbb{N}$ in (2.26) we derive

$$\begin{aligned}
 \|u_{n+1} - x_{n+1}\| &\leq [1 - A(1 - \delta)]\|u_n - x_n\| \\
 &\quad + [1 - A(1 - \delta)]\|u_n - Tu_n\| \\
 (2.28) \quad &\quad + [1 + \alpha_n\beta_n\delta]\varphi(\|u_n - Tu_n\|).
 \end{aligned}$$

Define

$$\begin{aligned}
 a_n &: = \|u_n - x_n\|, \\
 \mu_n &: = A(1 - \delta) \in (0, 1), \\
 \rho_n &: = [1 - A(1 - \delta)]\|u_n - Tu_n\| + [1 + \alpha_n\beta_n\delta]\varphi(\|u_n - Tu_n\|).
 \end{aligned}$$

Since $\lim_{n \rightarrow \infty} \|u_n - p\| = 0$, $\lim_{n \rightarrow \infty} \|u_n - Tu_n\| = 0$ as in the proof of Theorem 1. It therefore follows from the same argument that employed in the proof of Theorem 1 that $\lim_{n \rightarrow \infty} x_n = p$.

We will prove now that if the S-iteration converges, then the Mann iteration does too.

Using (1.4), (1.6), and (1.11) we have

$$\begin{aligned}
 \|x_{n+1} - u_{n+1}\| &= \|(1 - \alpha_n)(Tx_n - u_n) + \alpha_n(Ty_n - Tu_n)\| \\
 &\leq (1 - \alpha_n)\|Tx_n - u_n\| + \alpha_n\|Ty_n - Tu_n\| \\
 (2.29) \quad &\leq (1 - \alpha_n)\|Tx_n - u_n\| + \alpha_n\delta\|y_n - u_n\| + \alpha_n\varphi(\|y_n - Ty_n\|).
 \end{aligned}$$

We now have the following estimates

$$\begin{aligned}
 \|y_n - u_n\| &= \|(1 - \beta_n)x_n + \beta_nTx_n - u_n\| \\
 &= \|(1 - \beta_n)x_n + \beta_nTx_n - u_n - \beta_nu_n + \beta_nu_n\| \\
 (2.30) \quad &\leq (1 - \beta_n)\|x_n - u_n\| + \beta_n\|Tx_n - u_n\|,
 \end{aligned}$$

$$\begin{aligned}
(2.31) \quad \|Tx_n - u_n\| &= \|Tx_n - x_n + x_n - u_n\| \\
&\leq \|Tx_n - x_n\| + \|x_n - u_n\|.
\end{aligned}$$

Relations (2.29), (2.30), and (2.31) lead to

$$\begin{aligned}
(2.32) \quad \|x_{n+1} - u_{n+1}\| &\leq [1 - \alpha_n(1 - \delta)] \|x_n - u_n\| \\
&\quad + [1 - \alpha_n + \alpha_n \beta_n \delta] \|Tx_n - x_n\| + \alpha_n \varphi(\|y_n - Ty_n\|).
\end{aligned}$$

Since $\beta_n \in [0, 1]$ for all $n \in \mathbb{N}$,

$$(2.33) \quad \alpha_n \beta_n \delta < \alpha_n \delta.$$

Utilizing inequality (2.33) and the assumption $\alpha_n \geq A > 0, \forall n \in \mathbb{N}$ in (2.32) we get

$$\begin{aligned}
(2.34) \quad \|u_{n+1} - x_{n+1}\| &\leq [1 - A(1 - \delta)] \|x_n - u_n\| \\
&\quad + [1 - A(1 - \delta)] \|Tx_n - x_n\| + \alpha_n \varphi(\|y_n - Ty_n\|).
\end{aligned}$$

Now define

$$\begin{aligned}
a_n &:= \|x_n - u_n\|, \\
\mu_n &:= A(1 - \delta) \in (0, 1), \\
\rho_n &:= [1 - A(1 - \delta)] \|Tx_n - x_n\| + \alpha_n \varphi(\|y_n - Ty_n\|).
\end{aligned}$$

Since $\lim_{n \rightarrow \infty} \|x_n - p\| = 0$, $\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0$ as in the proof of Theorem 1.

Now we have

$$\begin{aligned}
(2.35) \quad 0 &\leq \|y_n - Ty_n\| \\
&\leq \|y_n - p\| + \|Tp - Ty_n\| \\
&\leq \|y_n - p\| + \delta \|p - y_n\| + \varphi(\|p - Tp\|) \\
&= (1 + \delta) \|y_n - p\| \\
&\leq (1 + \delta)(1 - \beta_n) \|x_n - p\| + (1 + \delta) \beta_n \|Tx_n - Tp\| \\
&\leq (1 + \delta)[1 - \beta_n + \beta_n] \|x_n - p\| + (1 + \delta) \beta_n \varphi(\|p - Tp\|) \\
&= (1 + \delta) \|x_n - p\| \rightarrow 0 \text{ as } n \rightarrow \infty,
\end{aligned}$$

that is, $\lim_{n \rightarrow \infty} \|y_n - Ty_n\| = 0$, therefore using the same argument as in the proof of Theorem 1, it can be shown that $\lim_{n \rightarrow \infty} u_n = p$. \square

As shown by Şoltuz and Grosan ([24], Theorem 3.1), in a real Banach space X , the Ishikawa iteration $\{x_n\}_{n=0}^{\infty}$ given by (1.7) converges to the fixed point of T , where $T : E \rightarrow E$ is a mapping satisfying condition (1.4).

In 2007, Şoltuz ([25], Corollary 2) proved that Krasnoselskij (1.7), Mann (1.6), Ishikawa (1.8), Noor (1.9) and multistep (1.10) iterations are equivalent for quasi-contractive operators in a normed space setting.

In 2011, Chugh and Kumar ([19], Corollary 3.2) proved that Picard (1.5), Mann (1.6), Ishikawa (1.8), new two step (1.12), Noor (1.9) and SP (1.13) iterations are equivalent for quasi-contractive operators in a Banach space setting.

From the argument used in the proofs of ([24], Theorem 3.1), ([25], Corollary 2) and ([19], Corollary 3.2) we can easily get the following corollary:

Corollary 1. *$T : E \rightarrow E$ be an operator satisfying condition (1.4) with $F_T \neq \emptyset$. If the initial point is the same for all iterations, $\alpha_n \geq A > 0, \forall n \in \mathbb{N}$, then the following are equivalent:*

- (1) *The Picard iteration (1.5) converges to $p \in F_T$;*

- (2) The Krasnoselskij iteration (1.7) converges to $p \in F_T$.
- (3) The Mann iteration (1.6) converges to $p \in F_T$;
- (4) The Ishikawa iteration (1.8) converges to $p \in F_T$;
- (5) The new two step iteration (1.12) converges to $p \in F_T$;
- (6) The Noor iteration (1.9) converges to $p \in F_T$;
- (7) The SP iteration (1.13) converges to $p \in F_T$;
- (8) The Multistep iteration (1.10) converges to $p \in F_T$;

Together with Theorem 1 and Theorem 2, Corollary 1 leads to the following corollary:

Corollary 2. $T : E \rightarrow E$ be an operator satisfying condition (1.4) with $F_T \neq \emptyset$. If the initial point is the same for all iterations, $\alpha_n \geq A > 0$, $\forall n \in \mathbb{N}$, then the following are equivalent:

- (1) The Picard iteration (1.5) converges to $p \in F_T$;
- (2) The Krasnoselskij iteration (1.7) converges to $p \in F_T$.
- (3) The Mann iteration (1.6) converges to $p \in F_T$;
- (4) The Ishikawa iteration (1.8) converges to $p \in F_T$;
- (5) The new two step iteration (1.12) converges to $p \in F_T$;
- (6) The Noor iteration (1.9) converges to $p \in F_T$;
- (7) The SP iteration (1.13) converges to $p \in F_T$;
- (8) The Multistep iteration (1.10) converges to $p \in F_T$;
- (9) The new multistep iteration (1.14) (or (1.15)) converges to $p \in F_T$;
- (10) The S -iteration (1.11) converges to $p \in F_T$.

Acknowledgement 1. This work is supported by Yildiz Technical University Scientific Research Projects Coordination Unit under project number BAPK 2012-07-03-DOP02.

REFERENCES

- [1] B.E. Rhoades, S.M. Şoltuz, "The equivalence of Mann iteration and Ishikawa iteration for non-Lipschitzian operators", Int. J. Math. Math. Sci., No. 42, 2645–2651 (2003)
- [2] B.E. Rhoades, S.M. Şoltuz, "The equivalence between the convergences of Ishikawa and Mann iterations for an asymptotically pseudocontractive map", J. Math. Anal. Appl., **283**(2003), no. 2, 681–688.
- [3] B.E. Rhoades, S.M. Şoltuz, "On the equivalence of Mann and Ishikawa iteration methods", Int. J. Math. Math. Sci., No. 7, 451–459 (2001).
- [4] B.E. Rhoades, S.M. Şoltuz, "The equivalence of Mann iteration and Ishikawa iteration for a Lipschitzian Ψ -uniformly pseudocontractive and Ψ -uniformly accretive maps", Tamkang J. Math., **35**(2004), No. 3, 235–245.
- [5] B.E. Rhoades, S.M. Şoltuz, "The equivalence between the convergences of Ishikawa and Mann iterations for an asymptotically nonexpansive in the intermediate sense and strongly successively pseudocontractive maps", J. Math. Anal. Appl., **289**(2004), No. 1, 266–278.
- [6] B.E. Rhoades, S.M. Şoltuz, "The equivalence of Mann and Ishikawa iteration for Ψ -uniformly pseudocontractive or Ψ -uniformly accretive maps", Int. J. Math. Math. Sci., **46**(2004), 2443–2452.
- [7] B.E. Rhoades, S.M. Şoltuz, "The equivalence between Mann-Ishikawa iterations and multi-step iteration", Nonlinear Analysis, **58**(2004), 219–228.
- [8] B.E. Rhoades, S.M. Şoltuz, "The equivalence of Mann and Ishikawa iteration dealing with strongly pseudocontractive or strongly accretive maps", Panamer. Math. J., **14**(2004), No. 4, 51–59.
- [9] C.O. Imoru, M.O. Olantiwo, "On the stability of Picard and Mann iteration processes", Carpathian Journal of Mathematics, **19**(2003), no.2, 155–160.

- [10] D.R. Sahu, "Applications of the S -iteration process to constrained minimization problems and split feasibility problems", Fixed Point Theory, **12**(2011), no. 1, 187-204.
- [11] F. Gürsoy, V. Karakaya, B.E. Rhoades, "Data dependence results of a new multistep and S -iterative schemes for contractive-like operators". (Communicated).
- [12] H. Akewe, "Strong convergence and stability of Jungck-multistep-SP iteration for generalized contractive-like inequality operators", Advances in Natural Science, **5**(2012), no. 3, 21-27.
- [13] J.O. Olaleru, H. Akewe, "The equivalence of Jungck-type iterations for generalized contractive-like operators in a Banach space", Fasciculi Mathematici, (2011), 47-61.
- [14] M.A. Noor, "New approximation schemes for general variational inequalities", Journal of Mathematical Analysis and Applications, **251**(2000), no.1, 217-229.
- [15] M.O. Osilike, A. Udomene, "Short proofs of stability results for fixed point iteration procedures for a class of contractive-type mappings", Indian Journal of Pure and Applied Mathematics, **30**(1999), no.12, 1229-1234.
- [16] Phuengrattana, Withunand, Suantai, Suthep, "On the rate of convergence of Mann, Ishikawa, Noor and SP iterations for continuous functions on an arbitrary interval", Journal of Computational and Applied Mathematics, **235**(2011), 3006-3014.
- [17] R.P. Agarwal, D. O'Regan, D.R. Sahu, "Fixed point theory for lipschitzian type-mappings with applications", Springer, ISBN: 978-0-387-75817-6, 2009.
- [18] R.P. Agarwal, D. O'Regan, D.R. Sahu, "Iterative construction of fixed points of nearly asymptotically nonexpansive mappings", J. Nonlinear Convex Anal., **8**(2007), no.1, 61-79.
- [19] R. Chugh, V. Kumar, "Strong convergence of SP iterative scheme for quasi-contractive operators in Banach spaces", International Journal of Computer Applications, **31**(2011), no.5.
- [20] S.S. Chang, Y.J. Cho, J.K. Kim, "The equivalence between the convergence of modified Picard, modified Mann, and modified Ishikawa iterations", Math. Comput. Modelling, **37**(2003), no. 9-10, 985-991.
- [21] S. Ishikawa, "Fixed points by a new iteration method", Proc. Amer. Math. Soc., **44**(1974), 147-150.
- [22] S.M. Şoltuz, "An equivalence between the convergences of Ishikawa, Mann and Picard iterations", Math. Commun., **8**(2003), no. 1, 15-22.
- [23] S.M. Şoltuz, "A remark concerning the paper: An equivalence between the convergences of Ishikawa, Mann and Picard iterations", Rev. Anal. Numer. Theor. Approx., **33**(2004), no. 1, 95-96.
- [24] S.M. Şoltuz, T. Grosan, "Data dependence for Ishikawa iteration when dealing with contractive like operators", Fixed Point Theory and Applications, **2008**(2008), Article ID 242916, 7 pages.
- [25] S.M. Şoltuz, "The equivalence between Krasnoselskij, Mann, Ishikawa, Noor and multistep iterations", Mathematical Communications, **12**(2007), 53-61.
- [26] S. Thianwan, "Common fixed points of new iterations for two asymptotically nonexpansive nonself mappings in a Banach space", J. Comput. Appl. Math., (2008), doi: 10.1016/j.cam.2008.05.051.
- [27] T. Zamfirescu, "Fix point theorems in metric spaces", Archiv der Mathematik, **23**(1972), no.1, 292-298.
- [28] V. Berinde, "Picard iteration converges faster than the Mann iteration in the class of quasi-contractive operators", Fixed Point Theory Appl. (2004), no. 2, 97-105.
- [29] V. Berinde, "On the convergence of the Ishikawa iteration in the class of quasi contractive operators", Acta Mathematica Universitatis Comenianae, **73**(2004), no.1, 119-126.
- [30] W.R. Mann, "Mean value methods in iterations", Proc. Amer. Math. Soc., **4**(1953), 506-510.
- [31] X. Weng, "Fixed point iteration for local strictly pseudocontractive mapping", Proc. Amer. Math. Soc., **113**(1991), 727-731.

DEPARTMENT OF MATHEMATICS, YILDIZ TECHNICAL UNIVERSITY, DAVUTPASA CAMPUS, ESENLER, 34220 ISTANBUL, TURKEY

E-mail address: `faikgursoy02@hotmail.com;fgursoy@yildiz.edu.tr`

URL: `http://www.yarbis.yildiz.edu.tr/fgursoy`

Current address: Department of Mathematical Engineering, Yildiz Technical University, Davutpasa Campus, Esenler, 34210 Istanbul

E-mail address: `vkkaya@yildiz.edu.tr;vkkaya@yahoo.com`

URL: `http://www.yarbis.yildiz.edu.tr/vkkaya`

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405-7106, USA

E-mail address: `rhoades@indiana.edu`

URL: `http://www.math.indiana.edu/people/profile.phtml?id=rhoades`